

### **AMENDMENTS TO THE CLAIMS**

The following listing of claims replaces all prior versions of claims in the application:

#### **Listing of Claims:**

1. (Currently amended) A method for detecting an unbound form of a first member of a binding pair, the method comprising the steps of:

- (a) providing a first particle bound to a second member;
- (b) reacting the first particle bound to the second member with a sample, wherein the second member binds to the unbound form of the first member in the sample thereby forming a first complex comprising the second member bound to the first particle and the first member;
- (c) providing into the sample a second particle bound to a third member, the third member being different from the second member and being capable of binding to the first member;
- (d) forming a second complex comprising the first particle, the second member, the first member, the third member and the second particle, wherein the third member bound to the second particle bound binds to the first member in the first complex and the first particle agglutinated to the second particle; and
- (e) detecting the formation of ~~agglutinated first and second particles of the~~ second complex by measuring an increase of the turbidity of the sample thereby detecting the unbound form of the first member in the sample.

2. (Original) The method of claim 1, wherein the third member is an antibody which specifically binds to the first member.

3. (Previously presented) The method of claim 1, wherein at least one of the first and second particle comprises latex.
4. (Cancelled)
5. (Original) The method of claim 1, wherein steps (a) through (d) are performed sequentially.
6. (Original) The method of claim 1, wherein steps (a) through (d) are performed simultaneously.
7. (Currently amended) The method of claim 1, wherein ~~the amount of the agglutinated first and second particles~~ detecting the formation of the second complex is quantitated.
8. (Previously presented) The method of claim 1, wherein the first member comprises protein S.
9. (Currently amended) The method of claim ~~8~~ 1, wherein the second member comprises C4b-binding protein (C4BP) or a fragment of C4BP that binds to protein S ~~thereof~~.
10. (Original) The method of claim 1, wherein the sample is selected from the group consisting of blood, plasma, serum, saliva, CSF, urine, culture media, a cell suspension, a buffer and an artificially prepared fluid containing the first member.
11. (Original) The method of claim 1, wherein the second member binds to the first member at a single binding site.
12. (Original) The method of claim 11, wherein the third member binds to the first member at a single binding site which is different from the single binding site to which the second member binds.
13. (Original) The method of claim 1, wherein step (b) is performed within 0 to about 180 seconds.

14. (Previously presented) The method of claim 1, wherein the third member and the second member are in a molar ratio of between about 2 and 20.
15. (Previously presented) The method of claim 1, wherein the third member and the second member are in a molar ratio of between about 5 and 10.
16. (Previously presented) The method of claim 1, wherein the third member is present in an amount that is higher than an amount of the free first member in the sample.
17. (Previously presented) The method of claim 1, wherein the third member and the free first member in the sample are in a molar ration of between about 10 and 40.
18. (Currently amended) A composition or a kit for detecting an unbound form of a first member of a binding pair in a sample, the binding pair comprising the first member and a second member, the composition or the kit comprising:
  - a first particle bound to the second member;
  - a second particle bound to a third member, the third member being different from the second member and capable of binding to the first member at a binding site different from the second member, and wherein the second member and the third member do not bind to each other;
  - wherein the first member comprises protein S and the second member comprises C4BP or a fragment of C4BP that binds to protein S thereof, and wherein the first and the second particles, when agglutinated, are capable of causing an increase of the turbidity of the sample.
19. (Cancelled)
20. (Currently amended) The composition or the kit of claim 18, wherein the third member is an antibody ~~and the second member is not an antibody.~~

21. (Cancelled)

22. (Previously presented) The composition or the kit of claim 21, wherein the third member binds to the first member at a single binding site which is different from the single binding site to which the second member binds.

23–31. (Cancelled)

32. (Currently amended) A method for diagnosing thrombophilia comprising performing the method of claim 8, and further comprising comparing the amount of the second complex formed ~~amount of the agglutinated first and second particles~~ to the amount of the second complex formed ~~of the agglutinated first and second particles~~ in a sample derived from an individual without thrombophilia.

33. (Cancelled)

34. (Previously presented) The composition of claim 18, wherein the size of the first particle and the second particle ranges from about 50 nm to about 1000 nm.

35. (Previously presented) The composition of claim 18, wherein the size of the first particle and the second particle are different.